




PIPA= Polyamides with Pyrrole and Imidazole as major building unites, novel gene control drug candidates, sequence specific binding to dsDNA

OPOB= Discovery tools, high quality one cyclic peptide immobilized on one gel-type bead + deconvolution technologies



Middle sized drugs such as peptide, designable & difficult to prepare, although HiPep established: Assembly, purification and QC.

PIPA blocks binding of transcription factors inhibiting gene expression, can control expression of gene.

Peptide Vehicle (THL, 55, 4091, 2014) DDS and DNA-modification alkylation of the target DNA

DNA visualization by labelled PIPAs

No-toxicities have been found, 3-4 weeks passed through urine (~100%)

Target diseases: not possible by small molecules or antibodies

A list of synthetic PIPAs for clinical applications published as reagents <http://hipep.jp/pdf/catalog/R002E.pdf>






Pioneer work by P. Dervan 1998 : *Science*, 282, 111, *Nature*, 391, 468.

<https://hipep.com/?p=2834>



AGP for regenerative medicine = peptides exhibiting angiogenic and cell-adhesion properties and their bioconjugates, longer lasting AGP for transplantation

PepTenChip® System: Biodetection technology with novel principles, disease-marker-independent, gives objective diagnostics



The 4 key technologies for PepTenChip® system have been completed: *de novo* designed labeled structured peptides as capture molecules; amorphous carbon, novel chip substrates; array technology; a detector for on-site use/maintenance free, PepTenCam. Most attempts for the protein-chip development have been failed and withdrew, due to the sensitivity and reproducibility by immobilization, instead solution assays such as ELISA were the mainstream. The PepTenChip® system does not involve detection of specific molecules in a 1:1 manner, but its detection principle is different (PAT: JP, US & EU).

Structure activity relationship had been elicited.

NGF and/or bFGF have been used to give certain effects, although infiltration has been observed. AGP showed no such effects, but the duration of actions has been relatively shorter and not enough for complete engraftment. Focusing on longer half-life time novel AGP has been developed (PAT.P).

Assay using human cells, human umbilical vein endothelial cells, with quantitative evaluation were accomplished.

Stability was tested in cell lysate and human serum.

Cyclic AGPs contribute directly to neovascularization, no linear fragments were generated in cell culture.

The structural changes of analyte are reflected in the fluorescence intensity changes of capture peptides in a dose dependent manner. This changes can be visualized as a pattern designated a "Protein Fingerprint". Datamining is carried out by statistical analyses : multivariate analyses. Discovery of undefined materials related to diseases can be possible. PepTenChip® can be repeatedly used (not disposable), Class I Medical device; noninvasive diagnostics; on-site "Point of Care". Objective results can be given not dependent on clinicians personal skill. Some clinical applications are underway: (1) Precancer diagnostics for early detection using gastric juice, classification of gastric precancerous lesions; (2) Multiple sclerosis (MS), of which cause as well as positive disease markers are unknown, discrimination in MS and related diseases using human spinal fluid for objective indicators, not relying the skills of clinicians.



PAT. US, JP, EU; *BBRC*, 310, 153-157, 2003;
Bioorganic & Medicinal Chem, DOI:10.1016/j.bmc.2020.115685;
Int. J. Peptide Res and Therapeutics, DOI 10.1007/s10989-022-10404-2.
<https://hipep.com/?p=1559>

<https://hipep.com/?p=781>

